



<b>Ecosystem Service</b>	<b>Aesthetics from interactions with nature</b>
<b>CICES class name</b>	Characteristics of living systems that enable aesthetic experiences
<b>CICES Section</b>	Cultural (Biotic)
<b>CICES Class code</b>	3.1.2.4

### Sample Indicators

Indicator values from			
Experiment or direct measurement		Survey	
Expert assessment		Statistical- or census data	
Model or GIS		Literature values	
Stakeholder participation		Not provided	

Table 1: Field Scale

Indicator	Unit	Indicator values from
<sup>[2]</sup> Presence of water bodies	Not provided	
<sup>[2]</sup> Presence of sublime features, e.g., mountains	Not provided	
<sup>[3]</sup> Functional diversity: Colour richness of flowers	# of colour groups visible to humans: white, yellow, purple, violet	
<sup>[3]</sup> Functional intensity: Average size of flowers or discernible sub-sets of inflorescences (of colour groups visible to humans)	cm	
<sup>[3]</sup> Functional stability: Average species richness of flowers within groups visible to humans during the flowering season	# of species	
<sup>[3]</sup> Overall species richness of flowers in colour groups visible to humans	# of species	



[3] Overall species richness of flowers	# of species	
[4] Abundance of large butterflies (species with median wingspan >5.4 cm)	Not provided	
[4] Abundance of birds that are either: colourful species or species that people attract to their homes with feeders	Not provided	
[4] Ant species richness as predictor of the abundance of birds, including those described above.	Not provided	

Table 2: Farm Scale

Indicator	Unit	Indicator values from
[2] Presence of water bodies	Not provided	
[2] Presence of sublime features, e.g., mountains	Not provided	
[5] Aesthetic landscape enhancement by a specific feature	poor-fair-good-excellent	
[6] Roadside variation: number of "land use patches" intersected by or adjacent to all roads and paths, except motorways and railways, divided by total road length	km <sup>-1</sup>	
[6] Landscape variation: length of land cover "edges" per hectare land surface	km * ha <sup>-1</sup>	
[6] Share of farmers surveyed that state that their farm should look well-tended	%	
[6] Share of farmers surveyed that consider open landscapes valuable landscape elements	%	

Table 3: Regional Scale

Indicator	Unit	Indicator values from
[1] Complexity: -Number of independently perceived visual elements in the scene -Visual richness, the degree of scene intricateness and "how much is going on." -The amount of information or the number of elements in the immediate environment -The promise of more information if one has more time to	not provided	



<p>observe from the specific point</p> <ul style="list-style-type: none"> <li>-The degree of simplicity versus complexity in the spatial structure</li> <li>-Presence of multiple elements with diverse forms elements at a given resolution</li> <li>-Diversity, richness and interspersions of landscape</li> <li>-The amount of diversity or variety in a scene, the engaging amount of information</li> <li>-The perceived degree of landscape variety (from not varied to varied)</li> <li>-Composition, distribution, organization and variation of landscape elements contributing to visual richness and diversity</li> </ul>		
<p><sup>[1]</sup> Diversity:</p> <ul style="list-style-type: none"> <li>-The degree of perceived visual variation among landscape elements</li> <li>-Visual diversity; the number and degree of image elements or different features</li> <li>-The diversity of landscape components "as the expression of vertical relationships between land use and abiotic features."</li> <li>-"Simply describes differences in nature, quality or aspect", also "the nature and relative size of the fields within the farm."</li> <li>-Composition, diversity, and relative abundance (evenness) of landscape cover types and land uses</li> </ul>	not provided	⊘
<p><sup>[1]</sup> Heterogeneity: grain size, visual compartmentalization and versatility within the landscape</p>	not provided	⊘
<p><sup>[1]</sup> Biodiversity: diversity of plants, insects or specific ecological groups relevant to scenic properties</p>	not provided	⊘
<p><sup>[1]</sup> Texture: The attribute of visual quality evaluated as smooth, medium or rough, or proportion of the landscape area covered by it</p>	not provided	⊘
<p><sup>[1]</sup> Pattern: presence of regularly repeated elements or clear patterns</p>	not provided	⊘
<p><sup>[1]</sup> Variety:</p> <ul style="list-style-type: none"> <li>-Scene as being varied or diverse in overall content; "diversity of colors, textures, shapes and masses, forms and spaces or other visible attributes that add a diversity or mixture of visual experiences."</li> </ul>	not provided	⊘
<p><sup>[1]</sup> Color diversity and contrast:</p> <ul style="list-style-type: none"> <li>-Variety of colors, chromatic diversity, visual contrast among available colors</li> </ul>	not provided	⊘

<p>[1] 3D complexity:</p> <ul style="list-style-type: none"> <li>-Heterogeneity in tree height and vertical vegetation layers</li> <li>-Visual grouping, density and structuring of vegetation, thinning intensity (managed ecosystems)</li> <li>-Presence of specific structural vegetation forms such as a tree, bush</li> <li>-Presence/absence &amp; diversity of man-made elements, either overall or as a modification to the landscape, sometimes as an undesirable factor</li> </ul>	not provided	⊘
<p>[1] Edge:</p> <ul style="list-style-type: none"> <li>-Presence, amount or density of distinct borders between areas</li> <li>-Presence of linear edge features such as hedgerows, walls, tree lines; visual properties of field margins</li> <li>-Edge condition</li> </ul>	not provided	⊘
<p>[1] Relief:</p> <ul style="list-style-type: none"> <li>-Topographic heterogeneity, variability in relief, non-uniform geomorphology, the contrast between flat and sloping</li> </ul>	not provided	⊘
<p>[1] Ephemera and seasonality:</p> <ul style="list-style-type: none"> <li>-Presence of elements and types of land use that change with seasons or overtime</li> <li>-Perception of seasonal change</li> </ul>	not provided	⊘
<p>[1] Time depth:</p> <ul style="list-style-type: none"> <li>-Visual evidence of historical continuity and diversity, sometimes as architectural variety and presence of landmarks</li> <li>-Level of succession (in woodlands)</li> </ul>	not provided	⊘
<p>[2] Presence of water bodies</p>	Not provided	⊘
<p>[2] Presence of sublime features, e.g., mountains</p>	Not provided	⊘
<p>[6] Roadside variation: number of land use patches intersected by or adjacent to all roads and paths, except motorways and railways, divided by total road length</p>	km <sup>-1</sup>	 , 
<p>[6] Landscape variation: length of land cover "edges" per hectare land surface</p>	km * ha <sup>-1</sup>	 , 
<p>[6] Share of farmers surveyed that state that their farm should look well-tended</p>	%	 , 

[6] Share of farmers surveyed that consider open landscapes valuable landscape elements	%	 , 
[7] Natural-aesthetical value: expert opinion/regional preferences	Not provided	
[7] Recreation potential: number of visitors	#	
[18] Average travel cost of tourists	€ * yr <sup>-1</sup>	
[8] Visibility of particularly beautiful spots (e.g., mountains, open water, forests, heterogeneous landscapes)	Index 0 - 100	
[14] Occurrence of protected areas, large forests, water bodies	Not provided	
[9] Open landscapes: Share of land under agricultural cultivation (keeping landscapes open through agriculture is seen as increasing aesthetic value)	%	
[9] Diversity of landscapes: Shannon index of land use	-	
[10] Number of residential properties in the direct vicinity of major rivers (number of properties is seen here as an indicator for aesthetic appreciation and inspiration)	#	
[11] Spatial mapping by stakeholders: stakeholders could place green stickers on a map to mark the supply hotspots of this ecosystem service. Red stickers were used to mark locations where the supply of this service is declining. Two different sizes of stickers were used to represent a radius of 0.75 km or 1 km, respectively	Index 0 - 5	
[12] Modelled landscape aesthetic value for a viewpoint: 360° panoramic photos of representative landscapes are created and assigned aesthetic scores [1-10] by stakeholders. The response is used to calibrate a regression model that relates landscape elements within the photos with the assigned aesthetic score. The following features are considered in the model: <ul style="list-style-type: none"> <li>- Landscape metrics (area-weighted mean patch area distribution [m<sup>2</sup>])</li> <li>- median radius of gyration distribution [m<sup>2</sup>]</li> <li>- modified Simpson's evenness index [-]</li> <li>- number of patches [#]</li> <li>- patch richness [-]</li> </ul>	-	 , 



<ul style="list-style-type: none"> <li>- range perimeter–area ratio distribution [-]</li> <li>- coefficient of variation of shape index distribution [-]</li> <li>- median of shape index distribution [-].</li> <li>- Land use classes (Settlement [0/1], Road [0/1], Forest [0/1], Water [0/1])</li> <li>- Viewshed in three distance zones (near zone 0–1.5 km, middle zone 1.5–10 km, far zone 10–50 km) [m<sup>2</sup>]</li> </ul>		
<p>[13] Recreation &amp; aesthetic values: values for land cover classes. The matrix by Burkhard et al., 2012 (DOI: 10.1016/j.ecolind.2011.06.019) was adapted and used in this study.</p>	Index 0-5	
<p>[15] Flower diversity: Plants Simpson's biodiversity index</p>	Not specified	
<p>[16] Visual quality index (VQI), based on 19 parameters (terrain ruggedness, presence of: waterfalls, wells and springs, area of standing water, length of flowing water, presence of the coast, habitat richness, area of woodland, presence of single large trees, number of plant species, hedgerow length, number of vegetation colours, area of human-influenced land, number of spot utilities/quarries, building area, road length, dry-stone walls length, presence of scheduled ancient monuments, presence of designated historic parks or gardens, presence of listed buildings)</p>	Index 0 - 1	 , 
<p>[17] Utility sum based on the following indicators:</p> <ul style="list-style-type: none"> <li>-Level of the presence of linear landscape elements within a grid cell [1 - 3]: hedgerows, tree rows, tree alleys and windbreaks</li> <li>-Level of the presence of point landscape elements within a grid cell [1 - 3]: hedgerows, tree rows, tree alleys and windbreaks</li> <li>-Level of presence of livestock within a grid cell [0 - 1]: occurrence of grasslands used as a proxy</li> <li>-Level of the diversity of crop production within a grid cell [1 - 3]: average plot size within field blocks used as a proxy</li> </ul>	-	
<p>[19] Landscape beauty index; Values per land use class based on:</p> <ul style="list-style-type: none"> <li>- a questionnaire-based photo survey on alpine landscapes</li> <li>- topographical visibility analysis (from DEM)</li> <li>- Shannon index of landscape diversity (Shannon index)</li> </ul> <p>Each of the three components was weighted equally.</p>	Not provided	 , 
<p>[20] Area providing an aesthetic and inspiring environment</p>	ha	 ,  , 

[21] Aesthetic value of landscapes: values from landscape preference studies	Not provided	
[22] Cumulative viewshed: visibility of green areas (such as farmland and forest) from residential land (using the visibility function in ArcGIS's Spatial Analyst)	#	
[23] Landscape aesthetics and landmark: Participatory mapping. Respondents in an online survey mark on map areas in their region where different cultural ecosystem services are supplied. Then, the proportion of markings in each of the investigated land cover classes is calculated. After that, values are calculated for subregions. The proportions are multiplied with the area extent of the respective land cover classes in the sub-region, and result for all land cover classes are summed up.	ha	
[18] Willingness to pay (WTP) for landscape preservation considering likely landscape changes	€	

Table 4: National Scale

Indicator	Unit	Indicator values from
[2] Presence of water bodies	Not provided	
[2] Presence of sublime features, e.g., mountains	Not provided	
[24] Shannon Diversity Index of landscapes	-	
[25] Number of visitors in agricultural areas	#	

Table 5: Multinational Scale

Indicator	Unit	Indicator values from
[2] Presence of water bodies	Not provided	
[2] Presence of sublime features, e.g., mountains	Not provided	



## References

No.	Citation
1	Dronova I (2017) Environmental heterogeneity as a bridge between ecosystem service and visual quality objectives in management, planning and design. <i>Landscape and Urban Planning</i> 163: 90-106. DOI: 10.1016/j.landurbplan.2017.03.005
2	Carvalho-Ribeiro S, Correia TP, Paracchini ML, Schupbach B, Sang AO, Vanderheyden V, Southern A, Jones P, Contreras B, O'Riordan T (2016) Assessing the ability of rural agrarian areas to provide cultural ecosystem services (CES): A multi scale social indicator framework (MSIF). <i>Land Use Policy</i> 53: 8-19. DOI: 10.1016/j.landusepol.2015.04.024
3	Kutt L, Lohmus K, Rammi IJ, Paal T, Paal J, Liira J (2016) The quality of flower-based ecosystem services in field margins and road verges from human and insect pollinator perspectives. <i>Ecological Indicators</i> 70: 409-419. DOI: 10.1016/j.ecolind.2016.06.009
4*	Peters VE, Campbell KU, Dienno G, García M, Leak E, Loyke C, Ogle M, Steinly B, Crist TO (2016) Ants and plants as indicators of biodiversity, ecosystem services, and conservation value in constructed grasslands. <i>Biodiversity and Conservation</i> 25(8): 1481-1501. DOI: 10.1007/s10531-016-1120-z
5	Fleming WM, Rivera JA, Miller A, Piccarello M (2014) Ecosystem services of traditional irrigation systems in northern New Mexico, USA. <i>International Journal of Biodiversity Science, Ecosystem Services and Management</i> 10(4): 343-350. DOI: 10.1080/21513732.2014.977953
6	Andersson E, Nykvist B, Malinga R, Jaramillo F, Lindborg R (2015) A social-ecological analysis of ecosystem services in two different farming systems. <i>Ambio</i> 44(1): 102-112. DOI: 10.1007/s13280-014-0603-y
7*	Fürst C, Frank S, Witt A, Koschke L, Makeschin F (2013) Assessment of the effects of forest land use strategies on the provision of ecosystem services at regional scale. <i>Journal of Environmental Management</i> 127: 96-116. DOI: 10.1016/j.jenvman.2012.09.020
8	Gret-Regamey A, Weibel B, Bagstad KJ, Ferrari M, Geneletti D, Klug H, Schirpke U, Tappeiner U (2014) On the Effects of Scale for Ecosystem Services Mapping. <i>Plos One</i> 9(12): e112601. DOI: 10.1371/journal.pone.0112601
9	Huber R, Lehmann B (2010) Economies of Scope in the Agricultural Provision of Ecosystem Services: An Application to a High Cost Production Region. <i>German Journal of Agricultural Economics</i> 59(2): 91-105.
10	Liu S, Crossman ND, Nolan M, Ghirmay H (2013) Bringing ecosystem services into integrated water resources management. <i>Journal of Environmental Management</i> 129: 92-102. DOI: 10.1016/j.jenvman.2013.06.047
11	Palomo I, Martin-Lopez B, Zorrilla-Miras P, Del Amo DG, Montes C (2014) Deliberative mapping of ecosystem services within and around Donana National Park (SW Spain) in relation to land use change. <i>Regional Environmental Change</i> 14(1): 237-251. DOI: 10.1007/s10113-013-0488-5
12	Schirpke U, Timmermann F, Tappeiner U, Tasser E (2016) Cultural ecosystem services of mountain regions: Modelling the aesthetic value. <i>Ecological Indicators</i> 69: 78-90. DOI: 10.1016/j.ecolind.2016.04.001

\* The ecosystem service discussed on this factsheet is not a focus of the cited paper



No.	Citation
13*	Zhang ZM, Gao JF, Fan XY, Lan Y, Zhao MS (2017) Response of ecosystem services to socioeconomic development in the Yangtze River Basin, China. <i>Ecological Indicators</i> 72: 481-493. DOI: 10.1016/j.ecolind.2016.08.035
14	Bastian O, Lupp G, Syrbe RU, Steinháúßer R (2013) Ecosystem services and energy crops - Spatial differentiation of risks. <i>Ekologia Bratislava</i> 32(1): 13-29. DOI: 10.2478/eko-2013-0002
15	Quétier F, Lavorel S, Daigney S, de Chazal J (2009) Assessing ecological and social uncertainty in the evaluation of land-use impacts on ecosystem services. <i>Journal of Land Use Science</i> 4(3): 173-199. DOI: 10.1080/17474230903036667
16	Swetnam RD, Harrison-Curran SK, Smith GR (2017) Quantifying visual landscape quality in rural Wales: A GIS-enabled method for extensive monitoring of a valued cultural ecosystem service. <i>Ecosystem Services</i> 26: 451-464. DOI: 10.1016/j.ecoser.2016.11.004
17	Ungaro F, Hafner K, Zasada I, Piorr A (2016) Mapping cultural ecosystem services: Connecting visual landscape quality to cost estimations for enhanced services provision. <i>Land Use Policy</i> 54: 399-412. DOI: 10.1016/j.landusepol.2016.02.007
18	van Berkel DB, Verburg PH (2014) Spatial quantification and valuation of cultural ecosystem services in an agricultural landscape. <i>Ecological Indicators</i> 37: 163-174. DOI: 10.1016/j.ecolind.2012.06.025
19	Vigl LE, Tasser E, Schirpke U, Tappeiner U (2017) Using land use/land cover trajectories to uncover ecosystem service patterns across the Alps. <i>Regional Environmental Change</i> 17(8): 2237-2250. DOI: 10.1007/s10113-017-1132-6
20	Adhikari S, Baral H, Nitschke CR (2018) Identification, Prioritization and Mapping of Ecosystem Services in the Panchase Mountain Ecological Region of Western Nepal. <i>Forests</i> 9(9): 554. DOI: 10.3390/f9090554
21	Duarte GT, Santos PM, Cornelissen TG, Ribeiro MC, Paglia AP (2018) The effects of landscape patterns on ecosystem services: meta-analyses of landscape services. <i>Landscape Ecology</i> 33(8): 1247-1257. DOI: 10.1007/s10980-018-0673-5
22	Hashimoto S, DasGupta R, Kabaya K, Matsui T, Haga C, Saito O, Takeuchi K (2018) Scenario analysis of land-use and ecosystem services of social-ecological landscapes: implications of alternative development pathways under declining population in the Noto Peninsula, Japan. <i>Sustainability Science</i> 14: 53-75. DOI: 10.1007/s11625-018-0626-6
23	Jaligot R, Chenal J, Bosch M, Hasler S (2019) Historical dynamics of ecosystem services and land management policies in Switzerland. <i>Ecological Indicators</i> 101: 81-90. DOI: 10.1016/j.ecolind.2019.01.007
24	Kirchner M, Schmidt J, Kindermann G, Kulmer V, Mitter H, Prettenthaler F, Rudisser J, Schuppenlehner T, Schonhart M, Strauss F, Tappeiner U, Tasser E, Schmid E (2015) Ecosystem services and economic development in Austrian agricultural landscapes - The impact of policy and climate change scenarios on trade-offs and synergies. <i>Ecological Economics</i> 109: 161-174. DOI: 10.1016/j.ecolecon.2014.11.005
25	Maes J, Liqueste C, Teller A, Erhard M, Paracchini ML, Barredo JI, Grizzetti B, Cardoso A, Somma F, Petersen JE, Meiner A, Gelabert ER, Zal N, Kristensen P, Bastrup-Birk A, Biala K, Piroddi C, Egho B, Degeorges P, Fiorina C, Santos-Martín F, Naruševičius V, Verboven J, Pereira HM, Bengtsson J, Gocheva K, Marta-Pedroso C, Snäll T, Estreguil C, San-Miguel-Ayanz J, Pérez-Soba M, Grêt-Regamey A, Lillebø AI, Malak DA, Condé S, Moen J, Czúcz B, Drakou



No.	Citation
	EG, Zulian G, Lavallo C (2016) An indicator framework for assessing ecosystem services in support of the EU Biodiversity Strategy to 2020. <i>Ecosystem Services</i> 17: 14-23. DOI: 10.1016/j.ecoser.2015.10.023